

A Climatology of Erythemal Ultraviolet Radiation, 1979–1992

**Sasha Madronich, Bernhard Mayer, and
Chris Fischer**

**NCAR, P.O. Box 3000, Boulder, CO 80307;
e-mail sasha@ucar.edu, bmayer@ucar.edu, fischer@ucar.edu**

**National
Center for
Atmospheric
Research**

Acknowledgments

This work was supported by the Department of Energy through grant DE–A 105–94ER61879 to NCAR. B.M. was supported by a research grant from the German Academic Exchange Service (DAAD). The National Center for Atmospheric Research is sponsored by the National Science Foundation.

1 Abstract

The distribution of erythematous (skin-reddening) ultraviolet radiation at the Earth's surface is derived from satellite-based observations of atmospheric ozone and cloud reflectivity. This climatology may be helpful in assessing the role of long-term environmental exposure to ultraviolet radiation, e.g., in epidemiological studies of skin cancer. Comparisons with ground-based measurements show agreement at the 10–20% level, except at high latitudes.

2 Methods

Total ozone and cloud/aerosol reflectivity at 380 nm were taken from the Total Ozone Mapping Spectrometer (Nimbus-7 TOMS, Level 3/V.7, [5]) from Jan. 1979 to Dec. 1992, at a geographical resolution of 1.25° longitude by 1.00° latitude. Total ozone was used as input to the Tropospheric Ultraviolet-Visible (TUV) model [3], the effect of clouds was considered in a second step [1]. Spectral irradiance $F(\lambda)$ at the Earth's surface was calculated every 30 minutes (1979-92) at 1 nm steps from 280 to 400 nm, using a 4-stream discrete ordinates method [7] with pseudo-spherical correction for improved accuracy at low sun [6]. Erythemally effective irradiance was calculated as

$$UV_{\text{ery}} = \int B(\lambda) F(\lambda) d\lambda \quad (1)$$

where $B(\lambda)$ is the CIE erythema sensitivity function (action spectrum) [4].




Figure 2: Daily erythemal UV doses averaged over 1979–92, United States.

3 Results and Discussion

Figure 1 shows the geographical distribution of daily UV_{ery} doses (top) and the corresponding cloudless sky values (bottom), averaged over the entire time period. As expected, highest values are generally seen in the tropics, up to ca. $7 \text{ kJ m}^{-2} \text{ day}^{-1}$ in the eastern Pacific and eastern Africa, while middle latitudes of both hemispheres show a general pole-ward decrease from about 5 to $1 \text{ kJ m}^{-2} \text{ day}^{-1}$. Local highs are associated with higher elevations, smaller ozone columns, and infrequent cloudiness (e.g., Andes mountains, Tibet plateau, Mexico and southwestern US). A more detailed climatology for the contiguous United States is shown in Figure 2.




Figure 1: Daily erythemal UV doses averaged over 1979–92. (Top) Including clouds, (Bottom) cloudless sky.

Comparisons with ground-based measurements, as archived at the World Ozone and UV Data Center (WOUDC) [8], are shown in Figure 3, for annually averaged daily UV_{ery} doses. With the exception of Palmer station, satellite-derived annual averages tend to overestimate measurements by ca. 10–20%, as also found by [2] for the Toronto measurements. These discrepancies remain under investigation and may stem from both measurements (e.g., cosine error) and satellite-derived values (e.g., local pollution) [2]. The disagreement at Palmer station is a consequence of using TOMS-observed reflectivity to infer cloudiness, when in reality the high reflectivity may be due to snow and ice, thus leading to a significant underestimate of UV_{ery} .




Figure 3: Annually averaged daily UV_{very} doses; satellite-derived climatology and ground-based measurements. The dashed lines are cloudless-sky calculations.

This nearly-global UV climatology may be a valuable tool in the study of the consequences of UV exposure in the biosphere, for example in epidemiological studies of human skin cancer, and in the assessment of long-term UV stresses on terrestrial and aquatic ecosystems.

References

- [1] Eck, T. F., P. K. Bhartia, and J. B. Kerr, *Geophys. Res. Lett.*, 22, 611–614, 1995.
- [2] Herman, J. R., N. Krotkov, E. Calarier, D. Larko, and G. Labow, *J. Geophys. Res.*, in press, 1999.
- [3] Madronich, S. and S. Flocke, Theoretical estimation of biologically effective UV radiation at the Earth's surface, in *Solar Ultraviolet Radiation – Modeling, Measurements and Effects* (Zerefos, C., ed.). NATO ASI Series Vol. I52, Springer–Verlag, Berlin, 1997.
- [4] McKinlay, A.F. and B. L. Diffey, A reference action spectrum for ultraviolet induced erythema in human skin. In: Passchier W.R. and Bosnjakovic, B.F.M. (eds.) *Human Exposure to Ultraviolet Radiation: Risks and Regulations*. Elsevier, Amsterdam, 1987.
- [5] McPeters, R. D. et al., Nimbus–7 Total Ozone Mapping Spectrometer (TOMS) Data Products User's Guide, NASA Reference Publication 1384, National Aeronautics and Space Administration, Washington, D.C. 1996.
- [6] Petropavlovskikh, I., Evaluation of photodissociation coefficient calculations for use in atmospheric chemical models, Ph. D. Thesis, U. of Brussels and NCAR/CT–159, Boulder, 1995.
- [7] Stamnes, K., S. Tsay, W. Wiscombe, and K. Jayaweera, *Appl. Opt.*, 27, 2502–2509, 1988.
- [8] WOUDC (World Ozone and UV Data Center), CD–ROM Volume 1, Environment Canada, Toronto, August 1996.